Nesting Characteristics of Quail Hollow Birds 2023*

SPECIES (list in order of # QH nests)	# QH Nests	Thumbnail Shots	Egg Appearance	Nest	# Eggs	# Broods	Incubation Days	Hatching to Fledging	Seasonal Nesting Range
Violet-green Swallow VGSW	127		White, no marks	Deep nest, grass or pine needles, feathers on top.	2-6 (3-8)	1 (1)	13-18 (13-14)	22-27 (20-21)	4/13-8/7 (4/15-
									7/31)
Chestnut- backed Chickadee CBCH	104		White to cream with rd/brn marks	Mossy lined with fur or plant down	3-8	1-2	12-16	18-23	3/6-6/24
					(5-8)	(1-?)	(11-12)	(1 3-17)	(3/15-
									6/20)
Western Bluebird WEBL	57		Blue, white, or mixed	Loose cup grass, stems	3-7	1-2	12-17	20-24	3/8-7/31
					(4-6)	(2-3)	(14)	(19-23)	(4/15-
									8/10)
Oak Titmouse OATI	36		White usually unmarked	Moss with grasses or pine needles, lined with fur	5-9	1	13-18	21-25	3/3-6/26
					(6-8)	(2)	(14-16)	(17-20)	(4/1-6/30)
Ash-throated Flycatcher ATFL	36		White to pink, splotched with browns and purples	Nest soft, of hair, fur feathers, grass	3-6	1-2	14-16	15-20	4/20-7/28
					(4-5)	(1)	(14-17)	(15-16)	(5/1-7/25)
Bewick's Wren BEWR	6		White w/ light spots of	Twigs, grass lined	5-7	1	14-15	15-17	4/23-6/11
		re	reds, greys and purples	with feathers and grass	(4-	(2)	(14)	(14)	(4/25-7/5)
					11)				
House Wren	4		White w/ tiny red	Twigs, cup in back	5-8	2	13-14	16-20	4/1-7/27
HOWR			brown dots on large end	lined with feathers	(5-8)	(2)	(12-15)	15-17	(4/10-7/5)

^{*} Quail Hollow numbers in **Bold.** Average numbers for California listed in parentheses taken from: Graham, Hatch. 2006. Monitoring Guide: Monitoring your Bluebird Trail in California. Hot Pepper Press. Somerset, CA.